Evolution of spherical cavitation bubbles: parametric and closed-form solutions

Stefan C. Mancas*
Hochschule München - Munich University of Applied Sciences, Germany

Haret C. Rosu[†]

IPICyT, Instituto Potosino de Investigacion Cientifica y Tecnologica, Camino a la presa San José 2055, Col. Lomas 4a Sección, 78216 San Luis Potosí, S.L.P., Mexico

We present an analysis of the Rayleigh-Plesset equation for a three dimensional vacuous bubble in water. In the simplest case when the effects of surface tension are neglected, the known parametric solutions for the radius and time evolution of the bubble in terms of a hypergeometric function are briefly reviewed. By including the surface tension, we show the connection between the Rayleigh-Plesset equation and Abel's equation, and obtain the parametric rational Weierstrass periodic solutions following the Abel route. In the same Abel approach, we also provide a discussion of the nonintegrable case of nonzero viscosity for which we perform a numerical integration.

PACS numbers: 47.55.dp, 02.30.Hq, 02.30.Ik

Keywords: Rayleigh-Plesset equation, cavitation, hypergeometric, Weierstrass, Emden-Fowler, Abel, Appell

invariant

INTRODUCTION

It is well established that the size evolution of unstable, spherical cavitation bubbles is governed by the Rayleigh-Plesset (RP) equation [1–3]

$$\rho_w \left(R\ddot{R} + \frac{3}{2}\dot{R}^2 \right) = p - P_\infty - \frac{2}{R} \left(\sigma + 2\mu_w \dot{R} \right) \,. \tag{1}$$

In (1) ρ_w is the density of the water, R(t) is the radius of the bubble, p and P_∞ are respectively the pressures inside the bubble and at large distance, σ is the surface tension of the bubble, and μ_w is the dynamic viscosity of water. In the simpler form with only the pressure difference in the right hand side, equation (1) was first derived by Rayleigh in 1917 [1] but it was only in 1949 that Plesset developed the full form of the equation and applied it to the problem of traveling cavitation bubbles [3]. In the second half of the last century a steady progress has been achieved with driving forces from engineering, medical, sonoluminescence, microfluidics, and even pharmaceutical applications of the cavitation phenomena [4]. In addition, the interest in the analytical and numerical solutions of cavitation dynamics remained considerable as these solutions can lead to a better control and understanding of the bubble collapse processes. In an effort to discern the peculiar features of the usual three-dimensional collapse, Prosperetti [2] and more recently Klotz [5] worked out generalizations to N-dimensional bubble dynamics.

In this paper, we first review the analytic solutions of the RP equation in terms of hypergeometric functions when the surface tension is neglected, and in terms of Weierstrass elliptic functions when the surface tension is taken into account [6, 7]. In the latter case, we employ an Abel equation approach which is a novel mathematical way of looking to the nonlinear evolution of cavitation bubbles. On the other hand, when the viscous term is introduced, we show that an Abel equation with a non constant invariant occurs. Since it is not yet known how to find analytical solutions to this equation (if any), we resort to numerical integration of the stiff RP equation from t = 0 to $t = t_c$, where t_c is the time of collapse of the bubble.

SIZE EVOLUTION AND COLLAPSE OF A SPHERICAL BUBBLE WITHOUT SURFACE TENSION: REVIEW OF CANONICAL RESULTS

We first consider the idealized case whereby the viscosity of the water is neglected, since $\mu_w \ll 1$, and we further discard the effect of surface tension σ . For this case, there are also recent analytical approximations in [8], which are further discussed in [9], but here we are concerned with the standard results.

Let us consider a vacuous p = 0 bubble of radius R which is surrounded by an infinite uniform incompressible fluid, such as water, that is at rest at infinity. We remark that 'infinity' in the present context refers to distance far enough

away from the initial position of the bubble, and we further assume that the pressure at infinity is constant, P_{∞} = const. Neglecting the body forces acting on the bubble, we have from equation (1)

$$2R\ddot{R} + 3\dot{R}^2 = -2\frac{P_{\infty}}{\rho_{w}} \ . \tag{2}$$

Since $R^2\dot{R}$ is an integrating factor of (1) consequently we obtain by one quadrature

$$R^3 \dot{R}^2 = -\frac{2}{3} \frac{P_{\infty}}{\rho_w} R^3 + \mathcal{C} .$$

Using the initial conditions $R(0) = R_0$ and $\dot{R}(0) = 0$ we find the integration constant to be

$$\mathcal{C} = \frac{2}{3} \frac{P_{\infty}}{\rho_w} R_0^3 ,$$

and hence, we obtain

$$\dot{R}^2 = \frac{2}{3} \frac{P_\infty}{\rho_w} \left[\left(\frac{R_0}{R} \right)^3 - 1 \right] . \tag{3}$$

Note that one can find a simple novel particular solution for R(t) by substituting (3) into (2) to obtain the Emden-Fowler equation

$$\ddot{R} = \mathcal{A}t^n R^m \tag{4}$$

with $A = -\frac{3\mathcal{C}}{2}$, n = 0, m = -4, and particular solution

$$R_p(t) = \sqrt[5]{\frac{25\mathcal{C}}{4}} \left(t + \frac{2R_0^{\frac{5}{2}}}{5\sqrt{\mathcal{C}}} \right)^{\frac{2}{5}} = \sqrt[5]{\frac{25}{6}} \frac{P_\infty R_0^3}{\rho_w} \left(t + \frac{\sqrt{6}}{5} \sqrt{\frac{\rho_w}{P_\infty}} R_0 \right)^{\frac{2}{5}} . \tag{5}$$

However, this solution is obtained under the assumption of a nonzero integrating factor [8] and therefore it fails to satisfy the second initial condition. Other solutions that do not satisfy the initial conditions have been found previously by Amore and Fernández [9]. As noticed in [10], equation (3) can be also viewed as a conservation law for the dynamics of the radius of the bubble, since its kinetic energy can be expressed as

$$2\pi\rho_w R^3 \dot{R}^2 = \frac{4}{3}\pi P_\infty (R_0^3 - R^3) \ . \tag{6}$$

To proceed with the integration of equation (3) we will use the set of transformations as given by Kudryashov [6], namely $R = S^{\epsilon}$, $dt = R^{\delta}d\tau$, where ϵ, δ are constants that depend on the dimension of the bubble, and S, τ are the new dependent and independent variables, respectively. Applying the transformations upon (3), we obtain the new dynamics in S and τ

$$S_{\tau}^{2} = \frac{2}{3} \frac{P_{\infty}}{\rho_{w}} \frac{1}{\varepsilon^{2}} (R_{0}^{3} S^{-3\varepsilon} - 1) S^{2+2\varepsilon\delta - 2\varepsilon} . \tag{7}$$

To find S, one sets $\varepsilon = \frac{1}{N}$, and $\delta = N + 1$, where N = 3 is the dimension of the bubble, which will in turn reduce (7) to the simpler equation

$$S_{\tau} = \sqrt{\frac{6P_{\infty}}{\rho_w}} S\sqrt{R_0^3 \ S - S^2} \ . \tag{8}$$

By integrating the above with $S(0) = R_0^3$ we obtain the rational solution

$$S(\tau) = \frac{R_0^3}{\mathcal{B}\tau^2 + 1} \,,\tag{9}$$

where for convenience we set $\mathcal{B} = \frac{9\mathfrak{C}}{4}R_0^3 = \frac{3}{2}\frac{P_\infty}{\rho_w}R_0^6$. Once we determine S, we can find the parametric solutions for the bubble radius $R(\tau)$ and evolution time of the bubble $t(\tau)$ [6]

$$R(\tau) = \frac{R_0}{(\mathcal{B}\tau^2 + 1)^{\frac{1}{3}}} ,$$

$$t(\tau) = R_0^4 \int_0^{\tau} \frac{d\xi}{(\mathcal{B}\xi^2 + 1)^{\frac{4}{3}}} .$$
(10)

The integral for the evolution of the time for bubble can be calculated analytically in terms of hypergeometric functions to give [6]

$$t(\tau) = \frac{R_0^4 \tau}{2} \left[\frac{3}{\sqrt[3]{\mathcal{B}\tau^2 + 1}} - {}_2F_1\left(\frac{1}{2}, \frac{1}{3}; \frac{3}{2}; -\mathcal{B}\tau^2\right) \right] = R_0^4 \tau_2 F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{3}{2}; -\mathcal{B}\tau^2\right) . \tag{11}$$

To achieve the time of collapse one needs to allow $\tau \to \infty$. This leads to $\lim_{\tau \to \infty} t(\tau) = 0.000908681$ if we use, in S.I. units, $[\rho_w] = 1000 \ kg/m^3$, $[R_0] = 10^{-2} \ m$ and $[P_\infty] = 101325 \ Pa$. We point out that the same asymptotic value for the collapse time has been also obtained, albeit using a different approach, by Obreschkow *et al.* [8] and it is also obtained from Eq. 14 below by direct integration. Once we solve for τ as a function of R from the first equation of (10), and substituting it into the second equation of (10) we obtain the closed-form solution

$$t(R) = R_0 \sqrt{\frac{2}{3} \frac{\rho_w}{P_\infty}} \sqrt{\left(\frac{R_0}{R}\right)^3 - 1} \, _2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{3}{2}; 1 - \left(\frac{R_0}{R}\right)^3\right)$$
 (12)

which is plotted as R(t) in Fig. 1.




FIG. 1: Radius of the bubble in the absence of surface tension according to equation (12) for $[\rho_w] = 1000 \ kg/m^3$, $[R_0] = 10^{-2} \ m$ and $[P_\infty] = 101325 \ Pa$.

Next, we will find the time for the total collapse t_c of the bubble by integration of equation (3), which yields

$$t_c = \frac{1}{R_0^{3/2}} \sqrt{\frac{3}{2}} \frac{\rho_w}{P_\infty} \int_0^{R_0} \frac{R^{3/2}}{\sqrt{1 - \left(\frac{R}{R_0}\right)^3}} dR .$$
 (13)

This is just the N=3 case of the general N-dimensional formula given in [5]. If we let $R=R_0\sin^{2/3}\theta$, where $\theta \in [0, \pi/2]$ then the integral (13) transforms to

$$t_c = \frac{2R_0}{3} \sqrt{\frac{3}{2} \frac{\rho_w}{P_\infty}} \int_0^{\pi/2} \sin^{2/3} \theta \, d\theta,$$

which by comparing with the integral relation for Beta function, namely,

$$B(m,n) = 2 \int_0^{\pi/2} \cos^{2m-1} \theta \sin^{2n-1} \theta d\theta$$

leads to

$$t_c = \frac{R_0}{3} \sqrt{\frac{3}{2} \frac{\rho_w}{P_\infty}} B\left(\frac{1}{2}, \frac{5}{6}\right) = R_0 \frac{\Gamma(\frac{5}{6})}{\Gamma(\frac{4}{3})} \sqrt{\frac{\pi \rho_w}{6P_\infty}} = 0.914681 \ R_0 \sqrt{\frac{\rho_w}{P_\infty}} \ . \tag{14}$$

This solution can be also obtained from (4) by using the parametric solution 2.3.1-2.2 from Polyanin's book [11]. If we insert in (14) the same S.I. numbers as used in the hypergeometric asymptotics, we obtain again $t_c = 0.000908681$ sec.

SURFACE TENSION INCLUDED VIA ABEL'S EQUATION

When the surface tension term added, equation (1) can be written in the form

$$\ddot{R} + \frac{3}{2R}\dot{R}^2 + \frac{K_1}{R} + \frac{K_2}{R^2} = 0 , \qquad (15)$$

where we define $K_1 = \frac{P_{\infty} - P}{\rho_w}$ and $K_2 = \frac{2\sigma}{\rho_w}$. Proceeding as in [12], we first show that solutions to a general second order ODE of type

$$\ddot{R} + f_2(R)\dot{R} + f_3(R) + f_1(R)\dot{R}^2 + f_0(R)\dot{R}^3 = 0$$
(16)

may be obtained via the solutions to Abel's equation (17) of the first kind (and vice-versa)

$$\frac{dy}{dR} = f_0(R) + f_1(R)y + f_2(R)y^2 + f_3(R)y^3$$
(17)

using the substitution

$$\dot{R} = \eta(R(t)) , \qquad (18)$$

which turns (16) into the Abel equation of the second kind in canonical form

$$\eta \dot{\eta} + f_3(R) + f_2(R)\eta + f_1(R)\eta^2 + f_0(R)\eta^3 = 0.$$
 (19)

Moreover, via the inverse transformation

$$\eta(R(t)) = \frac{1}{y(R(t))} \tag{20}$$

of the dependent variable, equation (19) becomes (17). The invariant of Abel's equation, see [13], can be written as

$$\Phi(R) = \frac{1}{3} \left(\frac{df_2}{dR} f_3 - f_2 \frac{df_3}{dR} - f_1 f_2 f_3 + \frac{2}{9} f_2^3 \right)$$
 (21)

and when is a constant is an indication that Abel's equation is integrable.

By identification of equation (15) with (16), we see that $f_1(R) = \frac{3}{2R}$, $f_2(R) = f_0(R) = 0$, and $f_3(R) = \frac{K_1}{R} + \frac{K_2}{R^2}$, and hence the Kamke invariant is $\Phi(R) = 0$, therefore Abel's equation (17) becomes the Bernoulli equation

$$\frac{dy}{dR} = f_1(R)y + f_3(R)y^3 \tag{22}$$

which, by one quadrature, has the solution

$$y(R) = \frac{\pm\sqrt{3}R^{\frac{3}{2}}}{\sqrt{3\mathcal{D} - 2K_1R^3 - 3K_2R^2}} \,. \tag{23}$$

By using equations (20) and (18), we obtain

$$\dot{R}^2 = \frac{3\mathcal{D} - 2K_1R^3 - 3K_2R^2}{3R^3} \tag{24}$$

and via the same initial conditions we obtain the integration constant

$$\mathcal{D} = \frac{R_0^2}{3} (3K_2 + 2K_1 R_0) ,$$

which gives

$$\dot{R}^2 = \frac{2K_1(R_0^3 - R^3) + 3K_2(R_0^2 - R^2)}{3R^3} \ . \tag{25}$$

Notice that when $\sigma=0\to K_2=0$, and $p=0\to K_1=\frac{P_\infty}{\rho_w}$, then the above becomes equation (3). The new energy with surface tension is

$$2\pi\rho_w R^3 \dot{R}^2 = \frac{4\pi}{3} \left[P_\infty (R_0^3 - R^3) + 3\sigma (R_0^2 - R^2) \right] . \tag{26}$$

Thus, at the time of collapse we have $\frac{4\pi}{3}R_0^2(P_\infty R_0 + 3\sigma) = 0.42443J$ for a surface tension $[\sigma] = 10^{-3} N/m$. To integrate equation (25), first let us write it in a more convenient way, as

$$\dot{R}^2 = \frac{a_3}{R^3} + \frac{a_1}{R} + a_0 \tag{27}$$

with coefficients defined as $a_3 = R_0^2 \left(K_2 + \frac{2K_1}{3}R_0\right)$, $a_1 = -K_2$, and $a_0 = -\frac{2K_1}{3}$, and we will use the same set of transformations, namely $R = S^{\epsilon}$, $dt = R^{\delta}d\tau$ which give in turn

$$S_{\tau}^{2} = \frac{S^{2+2\varepsilon\delta}}{\varepsilon^{2}} (a_{0}S^{-2\varepsilon} + a_{1}S^{-3\varepsilon} + a_{3}S^{-5\varepsilon}) , \qquad (28)$$

where now we set $\epsilon = -3/N = -1$, and $\delta = \frac{N+1}{2} = 2$. Thus, we obtain the Weierstrass elliptic equation

$$S_{\tau}^2 = a_0 + a_1 S + a_3 S^3 \tag{29}$$

which in standard form is

$$\wp_{\tau}^{2} = 4\wp^{3} - g_{2}\wp - g_{3} \tag{30}$$

via the linear substitution [14]

$$S(\tau) = \frac{4}{a_3} \wp(\tau + 29.709; g_2, g_3) . \tag{31}$$

The germs of the Weierstrass function g_2, g_3 are given by

$$g_2 = -\frac{a_1 a_3}{4} = \frac{K_2 R_0^2}{4} \left(K_2 + \frac{2K_1 R_0}{3} \right)$$

$$g_3 = -\frac{a_0 a_3^2}{16} = \frac{K_1 R_0^4}{24} \left(K_2 + \frac{2K_1 R_0}{3} \right)^2 .$$
(32)

Substituting K_1 and K_2 into the germs, the solution to equation (29) becomes

$$S(\tau) = \frac{6\rho_w}{R_0^2(P_{\infty}R_0 + 3\sigma)}\wp(\tau + 29.709; g_2, g_3) , \qquad (33)$$

where $g_2 = 3.37751 \cdot 10^{-11} \ m^8/sec^4$, $g_3 = 1.92645 \cdot 10^{-8} \ m^{12}/sec^6$, and the constant in the front of Weierstrass function from (33) takes the value of 59215.2 \sec^2/m^5 . Once S is known we can find the parametric solutions for the radius and time of the bubble with surface tension as

$$R(\tau) = \frac{1}{S(\tau)} = \frac{R_0^2(P_{\infty}R_0 + 3\sigma)}{6\rho_w} \frac{1}{\wp\left(\tau + 29.709; \frac{R_0^2\sigma}{3\rho_w^2}(P_{\infty}R_0 + 3\sigma), \frac{R_0^4P_{\infty}}{54\rho_w^3}(P_{\infty}R_0 + 3\sigma)^2\right)}$$

$$t(\tau) = \int_0^{\tau} \frac{d\xi}{S(\xi)^2} = \frac{R_0^4(P_{\infty}R_0 + 3\sigma)^2}{36\rho_w^2} \int_0^{\tau} \frac{d\xi}{\wp\left(\xi; \frac{R_0^2\sigma}{3\rho_w^2}(P_{\infty}R_0 + 3\sigma), \frac{R_0^4P_{\infty}}{54\rho_w^3}(P_{\infty}R_0 + 3\sigma)^2\right)^2}.$$
(34)

Related plots are presented in Fig. 2.




FIG. 2: Parametric solutions for time of evolution (top), radius of the bubble (middle), and radius vs time (bottom) from equation (34) when surface tension is present for $[\rho_w]=1000~kg/m^3, [R_0]=10^{-2}~m, [P_\infty]=101325~Pa$ and $[\sigma]=10^{-3}~N/m$.




FIG. 3: Comparison between the exact solutions without (dotted curve) and with surface tension (dashed curve). Notice that when we have surface tension the time of collapse is smaller.

THE RAYLEIGH-PLESSET EQUATION WITH VISCOSITY

When we include the viscosity, equation (1) reads

$$\ddot{R} + f_2(R)\dot{R} + f_1(R)\dot{R}^2 + f_3(R) = 0 , \qquad (35)$$

where $f_2(R) = \frac{K_3}{R^2}$, and $K_3 = \frac{4\mu_w}{\rho_w}$ is a constant, and $f_1(R), f_3(R)$ being the same functions as before. Thus, Abel's equation (17) becomes

$$\frac{dy}{dR} = f_1(R)y + f_2(R)y^2 + f_3(R)y^3 . {36}$$

Now, we also find the Kamke invariant according to equation (21) and we obtain

$$\Phi(R) = \frac{K_3[4K_3^2 - 9R(3K_2 + 5K_1R)]}{R^6} \ . \tag{37}$$

In terms of the physical variables of the system, the invariant is

$$\Phi(R) = \frac{2\mu_w [64\mu_w^2 - 9\rho_w R(6\sigma + 5P_\infty R)]}{27\rho_w^3 R^6}$$
(38)

and because is not a constant, we will try to reduce (36) using the Appell invariant instead.

First, we will eliminate the linear term via the transformation $y(R) = R^{\frac{3}{2}}z(R)$ to obtain the reduced Abel equation

$$\frac{dz}{dR} = h_2(R)z^2 + h_3(R)z^3 , (39)$$

where $h_2(R) = \frac{K_3}{\sqrt{R}}$, and $h_3(R) = (K_1R + K_2)R$.

According to the book of Kamke [13], for equations of the type (39) for which there is no constant invariant one should change the variables according to

$$z(R) = \hat{z}(\zeta(R)) ,$$

$$\zeta(R) = \int h_2(R)dR = 2K_3\sqrt{R} ,$$
(40)

which lead to the canonical form

$$\frac{d\hat{z}}{d\zeta} = \hat{z}^2 + \Psi(\zeta)\hat{z}^3 , \qquad (41)$$

where

$$\Psi(\zeta) = \frac{h_3(R(\zeta))}{h_2(R(\zeta))} = \zeta^3(b_3 + b_5\zeta^2)$$
(42)

is the Appell invariant, and the constants b_i are

$$b_{3} = \frac{K_{2}}{8K_{3}^{4}} = \frac{\sigma\rho_{w}^{3}}{2^{10}\mu_{w}^{4}},$$

$$b_{5} = \frac{K_{1}}{32K_{3}^{6}} = \frac{P_{\infty}\rho_{w}^{5}}{2^{17}\mu_{w}^{6}}.$$
(43)

Choosing a dynamic viscosity of water $[\mu_w] = 1.002 \ cP = 1.002 \cdot 10^{-3} \ kg/(m \cdot sec)$ the b_i constants take the values of $b_3 = 9.68789 \cdot 10^{14} \ sec^2/m^5$, while $b_5 = 7.63836 \cdot 10^{32} \ sec^4/m^{10}$. The units of ζ are $m^{\frac{5}{2}}/sec$. This Abel equation is not integrable through quadratures, but numerically we integrate the RP equation (35), from t = 0 to the point of stiffness which is the point in time where the bubble collapses, see the numerical solution on Fig. 4.




FIG. 4: Numerical solution for the RP equation (35) when surface tension and viscosity are both present for $[\rho_w] = 1000 \ kg/m^3$, $[R_0] = 10^{-2} \ m$, $[P_\infty] = 101325 \ Pa$, $[\sigma] = 10^{-3} \ N/m$ and $[\mu_w] = 1.002 \cdot 10^{-3} kg/(m \cdot sec)$.




FIG. 5: Comparison between the numerical solution (continuous curve) of Fig. 4 and the exact solutions of Fig. 3. As one can see, the bubbles collapse fastest when there is surface tension (dashed curve) followed by the no surface tension and no viscosity case (dotted curve). The slowest time is achieved when we have viscosity together with surface tension (continuous curve). Right panel is the zoomed in version of the figure in the left panel.

CONCLUSION

In this work, we have considered the RP equation for the size evolution of a bubble in water. In the first part of the paper, we have surveyed the standard results in the absence of surface tension but in a different way from those usually pursued in the literature. We have obtained the closed form hypergeometric solutions of Kudryashov and Sinelshchikov although in a different but equivalent form. From an Emden-Fowler form of the RP equation in this case we have also obtained a particular solution which however does not satisfy the second initial condition. In the presence of surface tension and viscosity, we have employed a new approach based on Abel's equation. When only the surface tension is included, we have obtained the known parametric rational Weierstrass solutions, whereas when viscosity is added, the corresponding Abel equation does not have a constant invariant, which explains the nonintegrability in this case. A numerical integration obtained from this nonintegrable Abel route is presented graphically.

Acknowledgment

This research was partially supported by internal funding from Embry-Riddle Aeronautical University. We thank the reviewers for their appropriate remarks which improved the quality of the paper.

- * Electronic address: mancass@erau.edu
- † Electronic address: hcr@ipicyt.edu.mx
- [1] Lord Rayleigh, "VIII. On the pressure developed in a liquid during the collapse of a spherical cavity", Philos. Mag. Ser. 6, **34**, 94 (1917).
- [2] A. Prosperetti, "Bubbles", Phys. Fluids 16, 1852 (2004).
- [3] M.S. Plesset, "The dynamics of cavitation bubbles", ASME J. Appl. Mech. 16, 228 (1949).
- [4] F. Lugli and F. Zerbetto, "An introduction to bubble dynamics", Phys. Chem. Chem. Phys. 9, 2447 (2007).
- [5] A.R. Klotz, "Bubble dynamics in N dimensions", Phys. Fluids 25, 082109 (2013).
- [6] N. A. Kudryashov and D. I. Sinelshchikov, "Analytical solutions for problems of bubble dynamics", Phys. Lett. A 379, 798 (2015).
- [7] N. A. Kudryashov and D. I. Sinelshchikov, "Analytical solutions of the Rayleigh equation for empty and gas-filled bubble", J. Phys. A: Math. Theor. 47, 405202 (2014).
- [8] D. Obreschkow, M. Bruderer, and M. Farhat, "Analytical approximations for the collapse of an empty spherical bubble", Phys. Rev. E 85, 066303 (2012).
- [9] P. Amore and F.M. Fernández, "Mathematical analysis of recent analytical approximations to the collapse of an empty spherical bubble", J. Chem. Phys. 138, 084511 (2013).
- [10] D. Obreschkow, P. Kobel, N. Dorsaz, A. de Bosset, C. Nicollier, and M. Farhat, "Cavitation bubble dynamics inside liquid drops in microgravity", Phys. Rev. Lett. 97, 094502 (2006).
- [11] A.D. Polyanin and V.F. Zaitsev, Handbook of Exact Solutions for Ordinary Differential Equations (CRC Press, Boca Raton, 1995).
- [12] S.C. Mancas and H.C. Rosu, "Integrable Abel equations and Vein's Abel equation", Math. Meth. Appl. Sci., doi: 10.1002/mma.3575., (2015).
- [13] E. Kamke, Differentialgleichungen: Lösungsmethoden und Lösungen (Chelsea, New York, 1959).
- [14] E.T. Whittaker and G.N. Watson, A Course of Modern Analysis (Cambridge Univ. Press, Cambridge, 1927).